

Mental health

Covid-19 (Britain): Psychosis cases rise in England as pandemic hits mental health

Saturday 23 October 2021, by [PIDD Helen](#) (Date first published: 18 October 2021).

29% rise in referrals for first suspected episode of psychosis between April 2019 and April 2021

Cases of psychosis have risen over the past two years in [England](#) as an increasing number of people experience hallucinations and delusional thinking amid the stresses of the Covid-19 pandemic.

There was a 29% increase in the number of people referred to mental health services for their first suspected episode of psychosis between April 2019 and April 2021, [NHS data](#) shows.

The rise continued throughout the spring, with 9,460 referred in May 2021, up 26% from 7,520 in May 2019.

The charity Rethink Mental Illness is urging the government to invest more in early intervention for psychosis to prevent further deterioration in people's mental health from which it could take them years to recover.

It says the statistics provide some of the first concrete evidence to indicate the significant levels of distress experienced across the population during the pandemic.

A [study earlier this month](#) found that anxiety and depression around the world increased dramatically in 2020, with an estimated 76m extra cases of anxiety and 53m extra cases of major depressive disorder than would have been expected had Covid not struck. Women and young people were disproportionately affected, the researchers said.

[Psychosis](#) can involve seeing or hearing things that other people do not (hallucinations) and developing beliefs that are not based on reality (delusions), which can be highly distressing. It can be a symptom of mental illness such as schizophrenia, bipolar disorder or severe depression, but psychosis can also be a one-off, potentially triggered by a traumatic experience, extreme stress or drug and alcohol misuse.

Despite the continued pressure on mental health services, Rethink Mental Illness is highlighting the importance of rapid access to treatment to prevent further episodes of psychosis and reduce people's risk of developing severe mental illness.

Nice guidelines for people experiencing a suspected first episode of psychosis state they should receive an assessment within two weeks. However, the charity fears that if the increase in referrals is sustained, more people will have to wait longer for vital treatment.

Brian Dow, the deputy chief executive of Rethink Mental Illness, said: “Psychosis can have a devastating impact on people’s lives. Swift access to treatment is vital to prevent further deterioration in people’s mental health which could take them years to recover from.

“These soaring numbers of suspected first episodes of psychosis are cause for alarm. We are now well beyond the first profound shocks of this crisis, and it’s deeply concerning that the number of referrals remains so high. As first presentations of psychosis typically occur in young adults, this steep rise raises additional concerns about the pressures the younger generation have faced during the pandemic.

“The pandemic has had a gamechanging effect on our mental health and it requires a revolutionary response. Dedicated additional funding for mental health and social care must go to frontline services to help meet the new demand, otherwise thousands of people could bear a catastrophic cost.”

A spokesperson for the Department of [Health](#) and Social Care said: “It is vital everyone can get the right support when they need it and we are delivering the fastest expansion in mental health services in NHS history, backed by an additional £2.3bn a year by 2023/24, benefiting hundreds of thousands more people.

“On top of this, we’ve invested an additional £500m this year to help people whose mental health has been particularly impacted by the pandemic. All [NHS](#) mental health providers have established 24/7 urgent helplines, which have answered around three million calls during the pandemic.”

‘My head told me to deal with hearing voices’

Tom Dunning, 30, has a diagnosis of borderline personality disorder, social anxiety disorder and PTSD.

“I was about 22 or 23 when I first started experiencing symptoms of psychosis. It was pretty much overnight having after finishing my degree that my head told me to deal with hearing voices. Hearing them was a daily occurrence and it pretty much made me feel scared of myself because I didn’t think anything was wrong so I couldn’t tell anyone how I felt.

“I had experienced bullying in my childhood and now I felt like my own mental health was bullying me by the voices telling me to do things. Everyone around me started to notice differences and knew something was wrong but I was scared to acknowledge that. I eventually saw my GP and it was the first time that I thought I needed support but it was also the first time that I knew it was something I could get through.”

Helen Pidd

This article was amended on 20 October 2021. Due to errors in the source data supplied to us by Rethink Mental Health, a previous version incorrectly stated that there had been a 75% increase in

the number of people referred to mental health services for their first suspected episode of psychosis between April 2019 and April 2021. This figure – and the related underlying data – has been corrected in the text and subheading.

P.S.

- The Guardian. Mon 18 Oct 2021 00.01 BST. Last modified on Wed 20 Oct 2021 15.51 BST:
<https://www.theguardian.com/society/2021/oct/18/psychosis-cases-soar-in-england-as-pandemic-hits-mental-health>

- Support the Guardian
Available for everyone, funded by readers

[Contribute](#)

[Subscribe](#)